

Name Answer Key
Physics
Period _____

Date _____
Gravity and Circles WS #8
Mrs. Nadworny

Universal Gravitation & Circular Motion REVIEW

1. A star with a mass of 2.13×10^{17} kg and planet Blue, which has a mass of 4.13×10^{22} kg, are separated by a distance of 4.8×10^{11} m.
- a. Calculate the gravitational attractive force between the star and Blue.

$$F_G = \frac{Gm_1m_2}{r^2} = \frac{(6.67 \times 10^{-11} \frac{N \cdot m^2}{kg^2})(2.13 \times 10^{17} kg)(4.13 \times 10^{22} kg)}{(4.8 \times 10^{11} m)^2} = 2.5 \times 10^6 N \text{ toward}$$

- b. Calculate the acceleration due to gravity on planet Blue if it has a radius of 3.26×10^6 meters.

$$g = \frac{Gm_1}{r^2} = \frac{(6.67 \times 10^{-11} \frac{N \cdot m^2}{kg^2})(4.13 \times 10^{22} kg)}{(3.26 \times 10^6 m)^2} = 0.259 \frac{m}{s^2} \text{ down}$$

- c. What would the weight of an astronaut ($m = 95$ kg) who travels to planet Blue?

$$F_{grav} = mg = (95kg)(0.259 \frac{m}{s^2}) = 25N \text{ down}$$

2. Two objects, m_1 and m_2 , are separated by a distance r . What happens to the gravitational force between them when the following changes are made?
- a. M_1 is 9 times larger and the distance between them triples.

$$F_g = \frac{Gmm}{r^2} = \frac{(1)(9)(1)}{(3)^2} = \text{no change}$$

- b. The distance is cut in fourth.

$$F_g = \frac{Gmm}{r^2} = \frac{(1)(1)(1)}{(\frac{1}{4})^2} = 16x \text{ larger}$$

- c. M_1 is doubled and m_2 is 5 times larger.

$$F_g = \frac{Gmm}{r^2} = \frac{(1)(2)(5)}{1^2} = 10x \text{ larger}$$

3. A student whirls a 19.96 g rubber stopper above their head on a string with a radius of 0.318 meters. The stopper completes 10 revolutions in 4.89 seconds. The force on the stopper is 1.04 newtons.

- a. Calculate the Period of stopper.

$$T = \frac{\text{time}}{\text{revolutions}} = \frac{4.89s}{10} = 0.489s$$

- b. Calculate the speed of the stopper.

$$v = \frac{2\pi r}{T} = \frac{2\pi(0.318m)}{0.489s} = 4.09 \frac{m}{s}$$

- c. Calculate the centripetal acceleration using the speed.

$$a_c = \frac{v^2}{r} = \frac{(4.09 \frac{m}{s})^2}{0.318m} = 52.6 \frac{m}{s^2} \text{ inward}$$

- d. Calculate the centripetal acceleration using the force.

$$a_c = \frac{F_c}{m} = \frac{1.04N}{0.01996kg} = 52.1 \frac{m}{s^2} \text{ inward}$$

4. Two rocks are floating in deep space far from the influences of other celestial bodies. One rock has a mass of 4.32 kg and the other has a mass of 8.71 kg. The attractive force between them is 0.03427 newtons. How far apart are they?

$$d = \sqrt{\frac{Gm_1m_2}{F_G}} = \sqrt{\frac{(6.67 \times 10^{-11} \frac{N \cdot m^2}{kg^2})(4.32kg)(8.71kg)}{0.03427N}} = .000271m = 2.71 \times 10^{-4}m$$

5. A car is driving around a circular racetrack of radius 86.0 meters. It experiences a centripetal acceleration of 17.3 m/s² inward. What is the speed at which the car is traveling?

$$v = \sqrt{a_c r} = \sqrt{(17.3 \frac{m}{s^2})(86.0m)} = 38.6m/s$$

6. An object of mass m is traveling around a circle of radius r with speed v .
a. What happens to the centripetal force if the mass is five times larger?

$$F = \frac{mv^2}{r} = \frac{(5)(1^2)}{1} = 5x \text{ larger}$$

- b. What happens to the centripetal force if the speed is doubled?

$$F = \frac{mv^2}{r} = \frac{(1)(2)^2}{1} = \text{quadrupled}$$

- c. What happens to the centripetal force if the radius is cut in quarters?

$$F = \frac{mv^2}{r} = \frac{(1)(1^2)}{(\frac{1}{4})} = 4x \text{ larger}$$

- d. What happens to the centripetal acceleration if the mass triples?

$$a = \frac{v^2}{r} = \text{nothing}$$

- e. What happens to the centripetal acceleration if the radius is tripled?

$$a = \frac{v^2}{r} = \frac{1^2}{3} = \frac{1}{3} \text{ the size}$$

7. A 2.6 kg stopper is twirled in a circle of radius 1.24 meters with a constant speed of 3.39 m/s. What is the tension in the rope?

$$F_c = ma_c = \frac{mv^2}{r} = \frac{2.6kg(3.39 \frac{m}{s})^2}{1.24m} = 24N \text{ inward}$$

8. What is the period of a ball being swung around in a circle of radius 6.71 meters at 6.4 m/s?

$$T = \frac{2\pi r}{v} = \frac{2\pi(6.71m)}{(6.4 \frac{m}{s})} = 6.6s$$